

Valuations in Nilpotent Minimum Logic

Pietro Codara

Dipartimento di Informatica
Università degli Studi di Milano
Milan, Italy
e-mail: codara@di.unimi.it

Diego Valota

Artificial Intelligence Research Institute (IIIA)
CSIC
Campus UAB, 08193 Bellaterra, Spain
e-mail: diego@iiia.csic.es

Abstract—The Euler characteristic can be defined as a special kind of valuation on finite distributive lattices. This work begins with some brief consideration on the rôle of the Euler characteristic on NM algebras, the algebraic counterpart of Nilpotent Minimum logic. Then, we introduce a new valuation, a modified version of the Euler characteristic we call *idempotent Euler characteristic*. We show that the new valuation encodes information about the formulæ in NM propositional logic.

Keywords—NM logic; NM algebra; NM logic; valuation; Euler characteristic

I. INTRODUCTION

Let L be a distributive lattice. A function $\nu: L \rightarrow \mathbb{R}$ is a *valuation* if it satisfies

$$\nu(x) + \nu(y) = \nu(x \vee y) + \nu(x \wedge y) \quad (1)$$

for all $x, y, z \in L$. Recall that an element $x \in L$ is *join-irreducible* if it is not the bottom element of L , and $x = y \vee z$ implies $x = y$ or $x = z$ for all $y, z \in L$. When L is finite, it turns out [19, Corollary 2] that any valuation ν is uniquely determined by its values on the join-irreducible elements of L , along with its value at the bottom element \perp of L .

A special kind of valuation, introduced by V. Klee and G.-C. Rota, is the Euler characteristic, defined as follows.

Definition 1.1 ([16, p. 120], [19, p. 36]). *The Euler characteristic of a finite distributive lattice L is the unique valuation $\chi: L \rightarrow \mathbb{R}$ such that $\chi(x) = 1$ for any join-irreducible element $x \in L$, and $\chi(\perp) = 0$.*

In [9], [10], the authors investigate the notion of Euler characteristic in a particular case of finite distributive lattice: Gödel algebras, the algebraic counterpart of the many-valued logic known as Gödel logic¹. Specifically, they consider the Lindenbaum algebra of Gödel logic over a finite set of variables and then they investigate the values assigned by the Euler characteristic to each equivalence class of formulæ. It turns out that the Euler characteristic encode logical information about the formulæ, but such information is classical, i.e. coincide with the analogous notion defined in classical propositional logic; namely, the Euler characteristic of a formula is the number of Boolean assignments which makes the formula true. Further, the authors generalize the notion of Euler characteristic to a family of new valuations, the many-valued versions of the Euler characteristic. The latter valuations are shown to be able to separate many-valued tautologies from non-tautologies.

¹For background on Gödel logic see, e.g., [15]. The characterization of Gödel algebra used in the cited papers is provided in [3], [8], [11].

In this paper we approach the same problem on a different many-valued logic, the Nilpotent Minimum logic NM. We will briefly investigate the logical meaning of the Euler characteristic on NM algebras, the algebraic counterpart of NM logic, showing that such valuation, as is, can not carry information about assignments making a formula classically true. In order to obtain such a result we will introduce a new valuation, a modified version of the Euler characteristic we call *idempotent Euler characteristic*, and prove that such valuation indeed is capable of capturing the desired information.

The NM logic is briefly presented in the next section. Section III contains our main results. In Section IV we spend a few word to describe a particular schematic extension of NM logic, known as the logic NM^{*}. We easily obtain, as a corollary of our main result, that the *idempotent Euler characteristic* on NM^{*} algebras plays exactly the same rôle as the Euler characteristic on Gödel algebras. We conclude our work with some consideration on possible further results.

II. THE LOGIC OF THE NILPOTENT MINIMUM

A *triangular norm* (also called *t-norm*; see [17]) is a binary, commutative, associative and monotonically non-decreasing operation on $[0, 1]^2$ that has 1 as unit element. The *Nilpotent Minimum t-norm* is a first example of a left-continuous but not continuous t-norm. It has been introduced by Fodor [13], and it is defined as

$$x \odot y = \begin{cases} \min\{x, y\} & \text{if } x + y > 1, \\ 0 & \text{otherwise.} \end{cases} \quad (2)$$

for every $x, y \in [0, 1]$.

Hence, the *Nilpotent Minimum propositional logic* (NM for short) lies in the hierarchy of extensions of the *Monoidal T-norm based Logic* (MTL), introduced in [12] by Esteva and Godo. The propositional language of MTL is built over the binary connectives $\odot, \wedge, \rightarrow$ and the constant \perp . Usually derived connectives are $x \leftrightarrow y = (x \rightarrow y) \odot (y \rightarrow x)$, $x \vee y = ((x \rightarrow y) \rightarrow y) \wedge ((y \rightarrow x) \rightarrow x)$, the negation $\neg x = x \rightarrow \perp$, and the constant $\top = \neg \perp$. We let $\varphi^2 = \varphi \odot \varphi$.

The WNM logic is obtained from MTL by adding the axiom:

$$\neg(x \odot y) \vee ((x \wedge y) \rightarrow (x \odot y)), \quad (\text{WNM})$$

while NM logic is given by WNM plus involutivity axiom:

$$\neg\neg x \rightarrow x. \quad (\text{INV})$$

The aforementioned Gödel logic can be obtained by adding the idempotency axiom to MTL logic. If we add the axiom $\neg(\neg x^2)^2 \leftrightarrow (\neg(\neg x)^2)^2$ to NM, we obtain its negation fixpoint-free version, called NM^- [14].

The following form of *local* deduction theorem holds in NM logic [2],

$$\varphi \vdash_{NM} \psi \text{ if and only if } \vdash_{NM} \varphi^2 \rightarrow \psi. \quad (3)$$

Hence, we say that NM logic *proves* ψ from φ , in symbols $\varphi \vdash_{NM} \psi$, when $\varphi^2 \rightarrow \psi$ is a theorem of NM logic.

The algebraic semantic of MTL is given by the variety of *MTL algebras* [12]. As Gödel algebras are exactly the prelinear Heyting algebras, NM algebras are the prelinear Nelson algebras [7]. Hence, NM logic is to *Nelson logic* (constructive logic with strong negation) as Gödel logic is to Intuitionistic logic.

The algebraic variety of NM algebras corresponding to NM logic has a nice property, that it is *locally finite* [18]. This means that finitely generated free algebras are finite. Hence, a combinatorial treatment of free n -generated algebras is feasible. Indeed, a characterization of free n -generated NM algebras based on partially ordered sets (posets for short) has been given in [4].

In the next section we introduce some algebraic and combinatorial notion that will be useful throughout the paper.

A. NM algebras

Abusing notation, in the following we identify logical connectives with their algebraic interpretations. An algebra $\mathbf{A} = \langle A, \wedge, \vee, \odot, \rightarrow, \perp, \top \rangle$ of type $(2, 2, 2, 2, 0, 0)$ is a WNM algebra if and only if $\langle A, \wedge, \vee, \perp, \top \rangle$ is a bounded lattice, with top \top and bottom \perp , $\langle A, \odot, \top \rangle$ is a commutative monoid, and it satisfies the *residuation* equation, $x \odot y \leq z$ if and only if $x \leq y \rightarrow z$, the *prelinearity* equation $(x \rightarrow y) \vee (y \rightarrow x) = \top$, the *weak nilpotent minimum* equation $\neg(x \odot y) \vee ((x \wedge y) \rightarrow (x \odot y)) = \top$. Therefore, WNM algebras are a class of involutive residuated lattices. When the lattice order is total, \mathbf{A} is called a *chain*. A WNM algebra that satisfies the *involutivity* equation $(x \rightarrow \perp) \rightarrow \perp = x$ is called *NM algebra*, while a *Gödel algebra* is a WNM algebra that satisfies *idempotency*, that is $x \odot x = x$. Negation $\neg x$ is usually defined by $x \rightarrow \perp$. An NM algebra satisfying $\neg(\neg x^2)^2 \leftrightarrow (\neg(\neg x)^2)^2 = \top$ is called a *NM⁻ algebra*. Given an element x of a NM algebra \mathbf{A} , we say that x is *negative* when $x < \neg x$, x is *positive* when $x > \neg x$. We call x a *negation fixpoint* when $x = \neg x$. Note that if \mathbf{A} has a negation fixpoint, then it is unique.

The variety \mathbb{NM} of NM algebras is generated by the *standard* NM algebra $[0, 1] = \langle [0, 1], \wedge^{[0,1]}, \vee^{[0,1]}, \odot^{[0,1]}, \rightarrow^{[0,1]}, 0, 1 \rangle$ where $\odot^{[0,1]}$ is the NM t-norm (2), $x \wedge^{[0,1]} y = \min\{x, y\}$, $x \vee^{[0,1]} y = \max\{x, y\}$ and

$$x \rightarrow^{[0,1]} y = \begin{cases} 1 & \text{if } x \leq y \\ \max\{\neg x, y\} & \text{otherwise.} \end{cases} \quad (4)$$

for every $x, y \in [0, 1]$.

By the subdirect representation theorem [6] and the fact that subdirectly irreducible MTL algebras are chains [12], every NM algebra \mathbf{A} is isomorphic to a subdirect product of a

family $(C_i)_{i \in I}$ of NM chains, for some index set I . When \mathbf{A} is finite and not trivial, then the family $(C_i)_{i \in I}$ of non trivial chains is essentially unique up to reordering of the finite index set I . Hence, there exist $\pi_i : \mathbf{A} \rightarrow C_i$ such that $\pi_i(a) = a_i$ for every $a \in \mathbf{A}$. We call a_i the i^{th} -*projection* of a . Then, we can display every element a in \mathbf{A} by means of its projections $(a_i)_{i \in I}$.

Since every finite NM chain $C = \langle C, \odot, \rightarrow, \vee, \wedge, \perp, \top \rangle$ is a subalgebra of $[0, 1]$, then by (2) and (4) and the fact that $\neg^{[0,1]}x := x \rightarrow^{[0,1]} 0$, we have

$$x \odot y = \begin{cases} \min(x, y) & x > \neg y; \\ \perp & x \leq \neg y. \end{cases} \quad (5)$$

$$x \rightarrow y = \begin{cases} \top & x \leq y; \\ \max(\neg x, y) & x > y. \end{cases} \quad (6)$$

for all $x, y \in C$.

Note that, given a NM chain C , every $x \in C$ is either positive, negative or a negation fixpoint.

Denote by FORM_n the set of all well-formed formulæ of NM logic whose propositional variables are contained in $\{x_1, \dots, x_n\}$. Let \mathbf{A} be a NM algebra, with $a_1, \dots, a_n \in A$, and let $\varphi \in \text{FORM}_n$. By $\varphi^A(a_1, \dots, a_n)$ we denote the element of A obtained by the evaluation of φ in A interpreting every x_i with the corresponding a_i , in particular $x_i^A = a_i$. With this notation a formula φ is a *tautology* of NM logic if and only if for every algebra $A \in \mathbb{NM}$ and for every $a_1, \dots, a_n \in A$, $\varphi^A(a_1, \dots, a_n) = \top^A$. Moreover, given two logical formulæ φ and ψ , we say that they are *logically equivalent* if and only if $(\varphi \leftrightarrow \psi)^A = \top^A$, for every $A \in \mathbb{NM}$. In symbols, $\varphi \equiv \psi$. Note that \equiv is an equivalence relation. The algebra whose elements are the equivalence classes of formulæ of NM logic with respect to \equiv is called the *Lindenbaum Algebra* of NM and its elements are denoted $[\varphi]_{\equiv}$. The free n -generated algebra NM_n in \mathbb{NM} is the Lindenbaum algebra of the logical formulæ over the first n variables. Since $[0, 1]$ is generic for \mathbb{NM} , then NM_n is isomorphic to the subalgebra of $[0, 1]^{[0,1]^n}$ generated by the projection functions $(a_1, \dots, a_n) \mapsto a_i$. It follows that there exists a map from equivalence classes of formulæ $[\varphi]_{\equiv}$ to real-valued functions $f : [0, 1]^n \rightarrow [0, 1]$.

Given a finite poset F and $S \subseteq F$, the *lower set* of S is $\downarrow S = \{x \in F \mid x \leq y \text{ for some } y \in S\}$, and the *upper set* of S is $\uparrow S = \{x \in F \mid x \geq y \text{ for some } y \in S\}$. A *forest* is a finite poset such that for every $x \in F$ the lower set $\downarrow \{x\}$ is a chain. A forest with a bottom element is called a *tree*, and its bottom element is called *root*.

Let \mathbf{A} be a finite NM algebra. A nonempty subset S of A is called a *filter* of \mathbf{A} when S is an upper set, and for all $x, y \in S$ then $x \odot y \in S$. Since S is finite, then it has a minimum element $\bigwedge_{x \in S} x$ (that is, S is principal). We call *generator* of S the minimum element of the filter S . A filter S of A is *prime* if $S \neq A$ and for all $x, y \in A$, $x \vee y \in S$ implies $x \in S$ or $y \in S$. Note that, for every prime filter S of \mathbf{A} , its generator is an idempotent join irreducible element of \mathbf{A} . We consider the reverse inclusion as a partial order between prime filters, that is $S \leq S'$ if and only if $S' \subseteq S$, for every couple of filters S and S' .

Proposition II.1 ([2]). *The set of prime filters of NM_n ordered by reverse inclusion is a forest.*

As a direct consequence of Proposition II.1, when S is generated by a minimal idempotent join irreducible elements of \mathbf{NM}_n , then S is the root of a tree in the forest of prime filters of \mathbf{NM}_n . In such case, following the classical terminology, we say that S is maximal (with respect to the inclusion among filters).

We conclude the Section with a simple Lemma ² that will be useful in the following.

Lemma II.2. *Let 2 and 3 be the two-elements and the three-elements NM chains, respectively. Then, given a finite NM algebra \mathbf{A} and a maximal prime filter \mathfrak{p} , the quotient \mathbf{A}/\mathfrak{p} is either isomorphic to 2, or isomorphic to 3.*

Proof: Let $(C_i)_{i \in I}$ be the subdirect representation of \mathbf{A} , and let $p \in \mathbf{A}$ be the join irreducible element that generates \mathfrak{p} . Note that since \mathfrak{p} is maximal and prime, then p is minimal and idempotent.

Since p is join irreducible then there exists only one $j \in I$ such that $p_j \neq \perp_j$. Moreover, $p_j > \neg p_j$, for else $p_j \odot p_j = \perp_j$, in contradiction with the idempotency of p . Finally, since p is minimal, p_j is the least positive element in C_j . Moreover, if C_j does not have a negation fixpoint f , $\neg p_j$ is the greatest negative element in C_j , otherwise f covers $\neg p_j$.

Denote with $\sim_{\mathfrak{p}}$ the congruence associated to \mathfrak{p} . By the above discussion, if C_j does not have a negation fixpoint then $\mathbf{NM}_n/\mathfrak{p}$ is isomorphic to the two element NM chain $[p]_{\sim_{\mathfrak{p}}} > [\neg p]_{\sim_{\mathfrak{p}}}$. Otherwise, if C_j has a negation fixpoint f then \mathbf{A}/\mathfrak{p} is isomorphic to the three element NM chain $[p]_{\sim_{\mathfrak{p}}} > [f]_{\sim_{\mathfrak{p}}} > [\neg p]_{\sim_{\mathfrak{p}}}$. ■

III. VALUATIONS IN NM LOGIC

Since \mathbf{NM}_n is a finite distributive lattice whose elements are formulæ in n variables, up to logical equivalence, we can extend the scope of valuations to formulæ, as follows.

Definition III.1. *Let $\nu : \mathbf{NM}_n \rightarrow \mathbb{R}$ be a valuation on the finite distributive lattice \mathbf{NM}_n . The valuation $\nu(\varphi)$ of a formula $\varphi \in \text{FORM}_n$ is the number $\nu([\varphi]_{\equiv})$.*

As mentioned in the introduction, one of the goals of [10] is the interpretation of the logical meaning of the Euler characteristic on Gödel algebras. In that specific case, it turns out that the Euler characteristic of a formula φ coincide with the number of Boolean assignments satisfying φ .

Turning now to the case of NM-algebras, we can hope that the Euler characteristic $\chi(\varphi)$ of a formula φ encodes information about assignments making φ true. At least, this should work for the join irreducible elements of \mathbf{NM}_n . But, unfortunately, this is not the case. Indeed, take, for instance, the formula

$$\alpha = (X \leftrightarrow \neg X)^2 \wedge X.$$

A straightforward verification shows that for every assignments $\mu : \text{FORM}_1 \rightarrow [0, 1]$, $\mu(\alpha) < 1$. Moreover $[\alpha]_{\equiv}$ is a

join irreducible element of \mathbf{NM}_1 . Indeed, one can check that for every formula $\psi \in \text{FORM}_1$ such that $[\psi]_{\equiv} \leq [\alpha]_{\equiv}$, either $[\psi]_{\equiv} = [\alpha]_{\equiv}$, or $[\psi]_{\equiv} = \perp$. Thus, $\chi(\alpha) = 1$. Compare with Fig. 1.



Fig. 1: \mathbf{NM}_1 is isomorphic to the product of the three depicted NM chains ([1]). The dashed line on the left is the generator, while the dotted line on the right is $[\alpha]_{\equiv}$.

Since the truth value of α is strictly lower than 1 under any assignment, but the Euler characteristic of α is greater than 0, we can not directly interpret χ as a measure of the number of classes of assignments making a formula true. We do not discuss further the rôle of Euler characteristic in NM logic here. Instead, we provide a new valuation that, as we will see later in this section, can be interpreted similarly to how the Euler characteristic has been interpreted in Gödel logic in [10].

Let us introduce such a valuation, slightly different from the Euler characteristic, defined as follows.

Definition III.2. *We define the idempotent Euler characteristic $\chi^+ : \mathbf{NM}_n \rightarrow \mathbb{R}$ as the valuation on \mathbf{NM}_n such that*

- 1) $\chi^+(\perp) = 0$;
- 2) for each join irreducible element $g \in \mathbf{NM}_n$,

$$\chi^+(g) = \begin{cases} 1 & \text{if } g \odot g = g, \\ 0 & \text{otherwise.} \end{cases}$$

Remark. Observe that, if g is a join irreducible element, but $g \odot g \neq g$, then $g \odot g = \perp$.

The following proposition highlights a fundamental property of this newly defined valuation. The name given to the valuation is due to such property.

Proposition III.3. *Fix $n \geq 1$. The idempotent Euler characteristic satisfies, for every $x \in \mathbf{NM}_n$,*

$$\chi^+(x \odot x) = \chi^+(x)$$

Proof: Let $x \in \mathbf{NM}_n$. Three cases are to be considered.

- 1) $x \odot x = x$.
- 2) $x \odot x = \perp$.
- 3) $x \odot x = y$, with $y \in \mathbf{NM}_n$, $y \neq x$, and $y \neq \perp$.

If 1) holds the proposition immediately follows. Suppose 2) holds. We need to prove that $\chi^+(x) = \chi^+(\perp) = 0$. First, observe that for every $y \in \mathbf{NM}_n$ such that $y \leq x$, we have $y \odot y \leq x \odot x$. Thus, $y \odot y = \perp$. Let $G = \{g_1, \dots, g_m\}$ be the poset of join irreducibles of \mathbf{NM}_n such that $g_i \leq x$. Note that $x = \bigvee_{i=1}^m g_i$. We proceed by induction on the structure of G . If $m = 1$, then x is a join irreducible (an atom of \mathbf{NM}_n), $G = \{x\}$, and $\chi^+(x) = 0$. Let $m \geq 2$, and suppose (inductive hypothesis) that the proposition holds for

²We thank the anonymous referee for pointing out that Lemma II.2 can be generalized to any NM-algebra, and not just to finite ones. This follows from the fact that the quotient by a maximal filter is a simple algebra and that up to isomorphism the only simple NM-algebras are 2 and 3.

every element $y = \bigvee_{g \in G'} g$, with $G' \subsetneq G$. Suppose x is not a join irreducible (otherwise, the result follows by Definition III.2). Say, without loss of generality, that g_m is maximal in G , and let $y = \bigvee_{i=1}^{m-1} g_i$. By Equation (1),

$$\chi^+(x) = \chi^+(g_m) + \chi^+(y) - \chi^+(g_m \wedge y)$$

By Definition III.2, $\chi^+(g_m) = 0$. Further, by inductive hypothesis, $\chi^+(y) = 0$. Let G' be the poset of join irreducible g of \mathbf{NM}_n such that $g \leq y$. Since g_m is join irreducible, and it is maximal in G , $y \not\leq g_m$. Thus, $G' \subsetneq G$. By inductive hypothesis, $\chi^+(g_m \wedge y) = 0$. We conclude $\chi^+(x) = 0$.

Suppose, finally, that 3) holds. Let $z = \neg x \wedge x$. By monotonicity of \odot , we obtain $z \odot z = \perp$. Thus, $\chi^+(z) = 0$. Moreover, $y \wedge z \leq z$, thus $(y \wedge z) \odot (y \wedge z) = \perp$. Therefore, $\chi^+(y \wedge z) = 0$. Using the subdirect representation, one can see that $x = y \vee z$. We obtain

$$\chi^+(x) = \chi^+(y) + \chi^+(z) - \chi^+(y \wedge z) = \chi^+(y),$$

and the proposition is proved. \blacksquare

We do not provide here an example of the values of the idempotent Euler characteristic on a free NM algebra, because of the dimension of such structures (\mathbf{NM}_1 has 48 elements). However, a clarifying example is depicted in Fig. 2, for the case of \mathbf{NM}_1^- .

Lemma III.4. *Fix integer $n \geq 1$, and let $x \in \mathbf{NM}_n$. Then, $\chi^+(x)$ equals the number of minimal idempotent join-irreducible elements $g \in \mathbf{NM}_n$ such that $g \leq x$.*

Proof: Let $x \in \mathbf{NM}_n$. If $x = \perp$ the Lemma trivially holds. Suppose $x \odot x = \perp$, with $x \neq \perp$. By Proposition III.3, $\chi^+(x) = 0$. Observe that for all $y \leq x$, $y \odot y \leq x \odot x$, and thus $y \odot y = \perp$. That is, no idempotent element, except \perp , is under x , as desired.

Suppose now $x \odot x \neq \perp$. Let F be the forest of all idempotent join irreducible elements $g \in \mathbf{NM}_n$ such that $g \leq x$. Since $x \odot x \neq \perp$, we have $F \neq \emptyset$. Recall that $x = \bigvee_{g \in F} g$. We proceed by induction on the structure of F . If F has only one element, then $F = \{x\}$. By Definition III.2, $\chi^+(x) = 1$, as desired.

Let now $|F| > 1$, let $l \in F$ be a maximal element of F , let $F^- = F \setminus \{l\}$, and let x^- be the join of the elements of F^- . Observe that $x = x^- \vee l$. Denote by M and M^- the number of minimal elements of F , and F^- , respectively.

If l is a minimal element of F , then $M = M^- + 1$. Let $l^- = l \wedge x^-$. One can check (for instance, using the subdirect representation), the l^- satisfies $l^- \odot l^- = \perp$. Thus, by Proposition III.3, $\chi^+(l^-) = 0$. By (1), using the inductive hypothesis, we have $\chi^+(x) = \chi^+(l) + \chi^+(x^-) - \chi^+(l^-) = 1 + M^- - 0 = M$, as desired.

If l is not a minimal element of F , then $M = M^-$. Let $l^- = l \wedge x^-$. Clearly, the forest of idempotent join irreducible elements under l forms a chain, we denote L . Moreover, one easily see that the forest of idempotent join irreducible elements under l^- is the chain $L \setminus \{l\}$. Thus, $\chi^+(l^-) = 1$. By (1), we have $\chi^+(x) = \chi^+(l) + \chi^+(x^-) - \chi^+(l^-) = 1 + M^- - 1 = M$, as desired. \blacksquare

Lemma III.5. *Fix $n \geq 1$, and let $\varphi \in \text{FORM}_n$. Let $O(\varphi, n)$ be the set of assignments $\mu : \text{FORM}_n \rightarrow \{0, \frac{1}{2}, 1\}$ such that $\mu(\varphi) = 1$. Then, there is a bijection between $O(\varphi, n)$ and the set of minimal idempotent join irreducible elements $g \in \mathbf{NM}_n$ such that $g \leq [\varphi]_{\equiv}$.*

Proof: Equipping $\{0, \frac{1}{2}, 1\}$ with the structure of an NM algebra, the resulting chain will be isomorphic to the three-element NM algebra $\mathbf{3}$.

Fix an assignment $\mu : \text{FORM}_n \rightarrow \{0, \frac{1}{2}, 1\}$. Then, there exists a unique homomorphism $h_\mu : \mathbf{NM}_n \rightarrow \mathbf{3}$ defined by

$$h_\mu([\varphi]_{\equiv}) = \mu(\varphi). \quad (7)$$

Conversely, for every $h : \mathbf{NM}_n \rightarrow \mathbf{3}$ we can define a unique assignment $\mu_h : \text{FORM}_n \rightarrow \{0, \frac{1}{2}, 1\}$ such that

$$\mu_h(\varphi) = h([\varphi]_{\equiv}). \quad (8)$$

This yields a bijection between assignments $\mu : \text{FORM}_n \rightarrow \{0, \frac{1}{2}, 1\}$ and NM homomorphisms $h : \mathbf{NM}_n \rightarrow \mathbf{3}$. In particular, consider that $\mu_h(\varphi) = 1$ if and only if $h_\mu([\varphi]_{\equiv}) = 1$. Moreover, $h_\mu^{-1}(1)$ is a prime filter \mathfrak{p}_{h_μ} in \mathbf{NM}_n .

By Lemma II.2 and the fact that h_μ is an NM algebra homomorphism, \mathfrak{p}_{h_μ} has to be maximal. Hence, for every $\mu \in O(\varphi, n)$ we can associate the minimal idempotent join irreducible element in \mathbf{NM}_n that generates \mathfrak{p}_{h_μ} .

Conversely, for every \mathfrak{p} maximal prime filter in \mathbf{NM}_n there exists an NM algebra homomorphism $h_{\mathfrak{p}} : \mathbf{NM}_n \rightarrow \mathbf{3}$, induced by the natural quotient map $\mathbf{NM}_n \rightarrow \mathbf{NM}_n/\mathfrak{p}$ composed with the embedding $\mathbf{NM}_n/\mathfrak{p} \rightarrow \mathbf{3}$ given by Lemma II.2. Thanks to the bijection established by (7) and (8), we are able to associate an assignment $\mu_{h_{\mathfrak{p}}}$ with every minimal idempotent join irreducible element p in \mathbf{NM}_n . And the Lemma is settled. \blacksquare

Combining Lemma III.4 and Lemma III.5 we can now state our main result.

Theorem III.6. *Fix an integer $n \geq 1$. For any formula $\varphi \in \text{FORM}_n$, the valuation $\chi^+(\varphi)$ equals the number of assignments $\mu : \text{FORM}_n \rightarrow \{0, \frac{1}{2}, 1\}$ such that $\mu(\varphi) = 1$.*

Remark. If φ is a tautology in NM logic, then $\chi^+(\varphi) = 3^n$.

IV. VALUATIONS IN NM⁻ LOGIC

As mentioned in Section II, \mathbf{NM}^- is the schematic extension of NM logic obtained adding the axiom $\neg(\neg x^2)^2 \leftrightarrow (\neg(\neg x)^2)^2$. On the algebraic side we have that an NM algebra is an \mathbf{NM}^- algebra if and only if it does not have a negation fixpoint. Since Definitions III.1 and III.2 easily apply to the \mathbf{NM}^- case, we can consider the idempotent Euler characteristic on free n -generated \mathbf{NM}^- algebras. As we will see later in this Section, the results we obtain in this case are interesting, although easy corollaries of the results obtained in the previous Section.

First of all, observe that Proposition III.3 and Lemma III.4 clearly hold on \mathbf{NM}_n^- algebras. Furthermore, we can easily adapt Lemma II.2 (and its proof) to \mathbf{NM}_n^- algebras, as follows.

Lemma IV.1. *Let $\mathbf{2}$ be the two-elements NM chain. Then, given a finite NM^- algebra \mathbf{A} and a maximal prime filter \mathfrak{p} , the quotient \mathbf{A}/\mathfrak{p} is isomorphic to $\mathbf{2}$.*

Appealing at the proof of Lemma III.5, given a maximal prime filter \mathfrak{p} , there exists an embedding from the quotient NM_n^-/\mathfrak{p} to the two-elements NM chain $\mathbf{2}$. Lemma III.5 thus takes the following form, in the NM^- case.

Lemma IV.2. *Fix $n \geq 1$, and let $\varphi \in \text{FORM}_n$. Let $O(\varphi, n)$ be the set of assignments $\mu : \text{FORM}_n \rightarrow \{0, 1\}$ such that $\mu(\varphi) = 1$. Then, there is a bijection between $O(\varphi, n)$ and the set of minimal idempotent join irreducible elements $g \in NM_n^-$ such that $g \leq [\varphi]_{\equiv}$.*

This fact, together with a revised version of Lemma III.4, allow us to restate our main theorem for NM^- logic.

Theorem IV.3. *Fix an integer $n \geq 1$. For any formula $\varphi \in \text{FORM}_n$, the valuation $\chi^+(\varphi)$ equals the number of assignments $\mu : \text{FORM}_n \rightarrow \{0, 1\}$ such that $\mu(\varphi) = 1$.*

Remark. If φ is a tautology in NM^- , then $\chi^+(\varphi) = 2^n$.

Example 1. Consider the subdirect representation of NM_1^- given in Fig. 1. Since the free 1-generated NM^- algebra is a subalgebra of NM_1 , we can obtain NM_1^- by removing the three elements NM chain (it is the only NM chain in the subdirect product with a negation fixpoint). Indeed, NM_1^- is obtained as a product of the two 1-generated four-elements NM chains. In Fig. 2 the order structure of NM_1^- has been labelled with the values given by the idempotent Euler characteristic.




Fig. 2: The order structure of NM_1^- . Elements are labelled with their idempotent Euler characteristic.

V. CONCLUSION, AND FURTHER WORK

Our brief discussion on the (classical) Euler characteristic lead to the conclusion that a proper logical meaning for such valuations does not follow the intuition of [10]. We think a deeper investigation deserve to be done.

Further research also has to be done in order to obtain more expressive valuations, generalizing the idempotent Euler characteristic. Indeed, as in the Gödel logic case, the study of k -valued extensions of NM logic seems to be a feasible task.

Finally, an approach similar to the one presented here can be applied to other logics lying in the same hierarchy of Gödel and NM logics. An example is NMG logic [22], the logic

of the ordinal sum of Gödel and NM standard chains. The study of the Euler characteristic, or some modified versions of such valuation, on NMG algebras is a natural prosecution of this work. In order to address the more difficult case given by WNM logic, a useful and clarifying intermediate step is the study of RDP logic [21]. Indeed, the structure of join irreducible elements of RDP logic has already been investigated in [5], while a poset representations of its free n -generated algebras has been provided in [20].

ACKNOWLEDGMENT

We thank Stefano Aguzzoli and Vincenzo Marra for many useful discussions on the topics of this work.

The authors were supported by the MIUR-FIRB research project PNCE - Probability theory of non-classical events. Valota acknowledges also partial support from a Marie Curie INdAM-COFUND Outgoing Fellowship. The research reported in this paper was carried out while Valota was a postdoc fellow of the Dipartimento di Scienze Teoriche e Applicate (Università dell'Insubria), supported by the PNCE research project.

REFERENCES

- [1] S. Aguzzoli, S. Bova, and B. Gerla, "Free Algebras and Functional Representation," in *Handbook of Mathematical Fuzzy Logic*, P. Cintula, P. Hájek, and C. Noguera, Eds. College Publications, 2011.
- [2] S. Aguzzoli, M. Busaniche, and V. Marra, "Spectral Duality for Finitely Generated Nilpotent Minimum Algebras, with Applications," *Journal of Logic and Computation*, vol. 17, no. 4, pp. 749–765, 2007.
- [3] S. Aguzzoli, O. D'Antona, and V. Marra, "Computing minimal axiomatizations in Gödel propositional logic," *Journal of Logic and Computation*, vol. 21, no. 5, pp. 791–812, 2011.
- [4] S. Aguzzoli and B. Gerla, "Normal Forms and Free Algebras for Some Extensions of MTL," *Fuzzy Sets and Systems*, vol. 159, no. 10, pp. 1131–1152, 2008.
- [5] S. Bova and D. Valota, "Finitely Generated RDP-Algebras: Spectral Duality, Finite Coproducts and Logical Properties," *Journal of Logic and Computation*, 2011.
- [6] S. Burris and H. Sankappanavar, *A Course in Universal Algebra*. Springer-Verlag, 1981.
- [7] M. Busaniche and R. Cignoli, "Constructive logic with strong negation as a substructural logic," *Journal of Logic and Computation*, vol. 20, no. 4, pp. 761–793, 2010.
- [8] P. Codara, O. M. D'Antona, and V. Marra, "An analysis of Ruspini partitions in Gödel logic," *Internat. J. Approx. Reason.*, vol. 50, no. 6, pp. 825–836, 2009.
- [9] —, "The Euler characteristic of a formula in Gödel logic," in *Proceedings of The International Symposium on Multiple-Valued Logic*, 2010, pp. 108–112.
- [10] —, "Valuations in Gödel logic, and the Euler characteristic," *Journal of Multiple-Valued Logic and Soft Computing*, vol. 19, no. 1-3, pp. 71–84, 2012.
- [11] O. M. D'Antona and V. Marra, "Computing coproducts of finitely presented Gödel algebras," *Ann. Pure Appl. Logic*, vol. 142, no. 1-3, pp. 202–211, 2006.
- [12] F. Esteva and L. Godo, "Monoidal t-Norm Based Logic: Towards a Logic for Left-Continuous t-Norms," *Fuzzy Sets and Systems*, vol. 124, no. 3, pp. 271–288, 2001.
- [13] J. Fodor, "Nilpotent Minimum and Related Connectives for Fuzzy Logic," in *Proceedings of FUZZ-IEEE'95*, 1995, pp. 2077–2082.
- [14] J. Gispert, "Axiomatic Extensions of the Nilpotent Minimum Logic," *Reports on Mathematical Logic*, vol. 37, pp. 113–123.
- [15] P. Hájek, *Metamathematics of fuzzy logic*, ser. Trends in Logic—Studia Logica Library. Dordrecht: Kluwer Academic Publishers, 1998, vol. 4.

- [16] V. Klee, "The Euler characteristic in combinatorial geometry," *Amer. Math. Monthly*, vol. 70, pp. 119–127, 1963.
- [17] E. P. Klement, R. Mesiar, and E. Pap, *Triangular Norms*. Dordrecht: Kluwer Academic Publishers, 2000.
- [18] C. Noguera, F. Esteva, and J. Gispert, "On Triangular Norm Based Axiomatic Extensions of the Weak Nilpotent Minimum logic," *Mathematical Logic Quarterly*, vol. 54, no. 4, pp. 387–409, 2008.
- [19] G.-C. Rota, "On the combinatorics of the Euler characteristic," in *Studies in Pure Mathematics (Presented to Richard Rado)*. London: Academic Press, 1971, pp. 221–233.
- [20] D. Valota, "Poset Representation for Free RDP-Algebras," in *Probability, Uncertainty and Rationality*, ser. CRM Series, H. Hosni and F. Montagna, Eds. Edizioni della Scuola Normale Superiore, Pisa 2010, vol. 10.
- [21] S. Wang, "A Fuzzy Logic for the Revised Drastic Product t-Norm," *Soft computing*, vol. 11, no. 6, pp. 585–590, 2007.
- [22] S. Wang, B. Wang, and D. Pei, "A Fuzzy Logic for an Ordinal Sum t-Norm," *Fuzzy Sets and Systems*, vol. 149, no. 2, pp. 297–307, 2005.